

## Prosser Animal Hospital

460 Wine Country Rd, Prosser, WA, 99350

**Phone:** (509) 786-1393

**Fax:** (509) 786-7084

**Email:** [contact@prosseranimalhospital.com](mailto:contact@prosseranimalhospital.com)

**Website:** [www.prosseranimalhospital.com](http://www.prosseranimalhospital.com)

# Neutering in Dogs

## Why should I have my dog neutered?

Neutering should be considered if you are keeping any male dog as a pet. Remember that Guide Dogs for the Blind, Hearing Dogs for Deaf People, and Dogs for the Disabled are routinely neutered, and this does not impair their ability to perform their duties.

## What are the advantages of neutering my male dog?

- Reduces the risk of benign prostatic hyperplasia and prostatitis
- Reduces the risk of hormone-related diseases such as perianal adenoma, and perianal hernias
- Eliminates the risk of testicular cancer, the second most common cancer in intact dogs
- Removes sexual urges, which usually decreases roaming behaviors
- Reduces certain types of aggression



## Is neutering performed for any other reason?

Neutering may be used in an attempt to treat certain forms of aggression; however, this may or may not be effective without concurrent training or in individual dogs. In older dogs, the operation may be performed to treat testicular tumors and some prostate gland conditions. It is also used to control hormonal (testosterone) dependent diseases such as perianal adenomas.

## What are the disadvantages?

Most of the perceived disadvantages are false. The most quoted of these are that the dog will become fat, lazy, and useless as a guardian. Obesity is probably the most commonly quoted disadvantage of neutering. In most cases, obesity is the result of overfeeding and not exercising enough. By regulating your dog's diet and caloric intake, you can prevent obesity in both neutered and intact males.

Some people are concerned that their dog will not grow to be as large. While testosterone does increase muscle mass, it actually causes growth plates to close sooner. Thus, early neutering often results in taller dogs.

**Neutering doesn't cause a change in personality, guarding instincts, intelligence, playfulness and affection.**


## When should the operation be performed?

Most veterinarians recommend neutering at around four to six months of age. However, neutering at an earlier age, which is a common practice at animal shelters, does not appear to be detrimental.

## Is there any alternative to surgery?

There have been recent advances in non-surgical neutering. These involve injection of a compound directly into the testicle. You should discuss this treatment with your veterinarian to determine if it is appropriate for your pet, but many veterinary facilities do not perform or encourage this kind of procedure.

## Are there any dangers associated with the operation?



Neutering is considered a major operation and requires general anesthesia. With any anesthetic the risk of serious complications, including death, is always present. However, with modern anesthetics and monitoring equipment, the risk of a complication is very low. It has been said that your pet has a greater chance of being injured in a car wreck than having an anesthetic or surgical complication.

## What happens when my dog undergoes this procedure?

Your pet will be examined by a veterinarian and pre-anesthetic blood tests will usually be performed. If everything is acceptable, your pet will be anesthetized. After your pet is anesthetized, a breathing tube will be placed in his trachea or "windpipe" to deliver oxygen and gas anesthetic directly into the lungs. The surgery consists of making a small incision in front of the scrotum and removing the testicles. Many veterinarians use absorbable internal sutures so that you do not have to return your dog to the hospital to have them removed.

Used Under License  
Copyright© Lifelearn Inc.

## should take?

## Are there any post-operative precautions I

"Rest and restriction of activity are the primary post-operative care you should provide."

Rest and restriction of activity are the primary post-operative care you should provide. Most dogs can resume normal activity five to ten days after surgery. Until then, leash walks, lots of rest, and no swimming, bathing, running or climbing stairs are the rule. Some dogs will lick the incision excessively, which can not only traumatize the skin but introduce infection. Licking should be discouraged, and some dogs require an Elizabethan collar (or "cone" collar) to prevent licking.

*This client information sheet is based on material written by: Ernest Ward, DVM*

